

Microscopic locomotion and controllability of affine systems

Clément Moreau



Laboratoire des Sciences
du Numérique de Nantes


Séminaire d'automatique du plateau de Saclay — November 4th, 2025

Microscopic locomotors


Euglena gracilis



Volvox




Chlamydomonas



Mag. 1000X


Paramecium



Spermatozoon




Caenorhabditis elegans




Micro-robots


Sovolev et al., Small 5(14), 2009




Xin et al., ACS Nano 15(11), 2021





Ghosh & Fischer, Nano Letters 9(6), 2009




Magdanz et al., Adv. Funct. Mater. 25(18), 2015



^b Palagi, Fischer, Nat. Rev. Materials 3 (2018) ^c





ACS NANO

www.acsnano.org


Open Access

This article is licensed under [CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)  

Technology Roadmap of Micro/Nanorobots

REVIEW

Principles of micro-swimming





hydrodynamics

deformation

locomotion


Principles of micro-swimming



rigid motion

$\mathcal{X}(t)$ Pose in $SO(n)$

Principles of micro-swimming



hydrodynamics

deformation

locomotion

rigid motion

$\mathcal{X}(t)$


Pose in $SO(n)$

rate of deformation

$\dot{\mathcal{S}}(t)$

« Degrees of freedom », finite (or infinite) dimension

Principles of micro-swimming



hydrodynamics

deformation

locomotion

rigid motion

rate of deformation

Stokes flow

(Zero Reynolds number)


$\mathcal{X}(t)$ Pose in $SO(n)$

$\dot{\mathcal{S}}(t)$ « Degrees of freedom », finite (or infinite) dimension

$$\Delta \mathbf{u} + \nabla p = 0, \quad \nabla \cdot \mathbf{u} = 0$$

$$\text{B.C.} : \dot{\mathcal{X}} + \dot{\mathcal{S}}$$

Principles of micro-swimming



hydrodynamics

deformation

locomotion

rigid motion

$\mathcal{X}(t)$ Pose in $SO(n)$

rate of deformation

$\dot{\mathcal{S}}(t)$ « Degrees of freedom », finite (or infinite) dimension


Stokes flow

▪ Linearity:
 $\mathbf{F}^{\text{hydro}} = \mathbf{Q}(\mathcal{S})(\dot{\mathcal{X}}, \dot{\mathcal{S}})^T$

Stokes flow

$$\Delta \mathbf{u} + \nabla p = 0, \quad \nabla \cdot \mathbf{u} = 0$$

Principles of micro-swimming



hydrodynamics

deformation

locomotion

rigid motion

$\mathcal{X}(t)$ Pose in $SO(n)$

rate of deformation

$\dot{\mathcal{S}}(t)$ « Degrees of freedom », finite (or infinite) dimension

Stokes flow

Stokes flow

$$\Delta \mathbf{u} + \nabla p = 0, \quad \nabla \cdot \mathbf{u} = 0$$


▪ Linearity:

$$\mathbf{F}^{\text{hydro}} = \mathbf{Q}(\mathcal{S})(\dot{\mathcal{X}}, \dot{\mathcal{S}})^T$$

▪ Inertialess motion:

$$\mathbf{F}^{\text{hydro}} = 0$$

Principles of micro-swimming



hydrodynamics

deformation

locomotion

rigid motion

$\mathcal{X}(t)$ Pose in $SO(n)$

rate of deformation

$\dot{\mathcal{S}}(t)$ « Degrees of freedom », finite (or infinite) dimension

Stokes flow

Stokes flow

$$\Delta \mathbf{u} + \nabla p = 0, \quad \nabla \cdot \mathbf{u} = 0$$

▪ Linearity:

$$\mathbf{F}^{\text{hydro}} = \mathbf{Q}(\mathcal{S})(\dot{\mathcal{X}}, \dot{\mathcal{S}})^T$$

▪ Inertialess motion:

$$\mathbf{F}^{\text{hydro}} = 0$$

« Stokes connection »

$$\dot{\mathcal{X}} = \mathbf{N}(\mathcal{S})\dot{\mathcal{S}}$$


Swimming equation

Principles of microswimming


Consequence: Scallop theorem

"A swimmer with only one degree of freedom cannot produce net locomotion at low Reynold's number"


→ **Non-reciprocal** deformation is required



■ The famous **Purcell swimmer**





Purcell, Am. J. Phys., 1977




Three rigid links
with flexible junctions

Reciprocal gait




No net motion

Non-reciprocal gait



Net motion

“Non-reciprocal deformation is required for net locomotion”




Question:

Which kind of net locomotion?

Is **every direction** reachable as soon as I can do a “nonreciprocal” deformation?


→ **controllability**

Swimming is a driftless control-affine system



$$\dot{\mathcal{X}} = \mathbf{N}(\mathcal{S})\dot{\mathcal{S}}$$


Swimming equation



$$\dot{x} = \sum_{i=1}^m f_i(x)u_i$$


Control-affine system

Swimming is a driftless control-affine system



$$\begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$


Swimming equation



$$\dot{x} = \sum_{i=1}^m f_i(x) u_i$$


Control-affine system

Controllability




Control-affine system

Controllability: it is possible to go from x_0 to x_1 (for all x_0, x_1)




Controllability



Control-affine system

Local controllability: it is possible to go from x_0 to x_1 in a neighbourhood of an equilibrium



STLC: in small time and with small controls


B-STLC: in small time and with bounded controls

Lie brackets: formal non-reciprocity

$$\dot{x} = \sum_{i=1}^m f_i(x) u_i$$

$\searrow \quad m = 2$

$$\dot{x} = f_1(x) u_1 + f_2(x) u_2$$




Lie brackets: formal non-reciprocity

$$\dot{x} = \sum_{i=1}^m f_i(x) u_i$$

$\searrow \quad m = 2$

$$\dot{x} = f_1(x)u_1 + f_2(x)u_2$$




Lie brackets: formal non-reciprocity

$$\dot{x} = \sum_{i=1}^m f_i(x) u_i$$

$\searrow \quad m = 2$

$$\dot{x} = f_1(x)u_1 + f_2(x)u_2$$




Lie brackets: formal non-reciprocity

$$\dot{x} = \sum_{i=1}^m f_i(x) u_i$$

$\searrow \quad m = 2$

$$\dot{x} = f_1(x)u_1 + f_2(x)u_2$$




Lie brackets: formal non-reciprocity

$$\dot{x} = \sum_{i=1}^m f_i(x) u_i$$

$\searrow \quad m = 2$

$$\dot{x} = f_1(x)u_1 + f_2(x)u_2$$




Lie brackets: formal non-reciprocity

$$\dot{x} = \sum_{i=1}^m f_i(x) u_i$$

$\searrow \quad m = 2$

$$\dot{x} = f_1(x)u_1 + f_2(x)u_2$$





Lie brackets: formal non-reciprocity

$$\dot{x} = \sum_{i=1}^m f_i(x) u_i$$

$m = 2$

$$\dot{x} = f_1(x)u_1 + f_2(x)u_2$$



Lie brackets: formal non-reciprocity


$$\dot{x} = \sum_{i=1}^m f_i(x) u_i$$

$\searrow \quad m = 2$

$$\dot{x} = f_1(x)u_1 + f_2(x)u_2$$

$$x(4\varepsilon) = x_0 + \varepsilon^2[f_1, f_2](x_0) + o(\varepsilon^2)$$

$$[f_1, f_2](x) = \nabla f_2(x)f_1(x) - \nabla f_1(x)f_2(x)$$



Lie brackets: formal non-reciprocity


$$\dot{x} = \sum_{i=1}^m f_i(x) u_i$$

$m = 2$

$$\dot{x} = f_1(x) u_1 + f_2(x) u_2$$

$$x(4\varepsilon) = x_0 + \varepsilon^2 [f_1, f_2](x_0) + o(\varepsilon^2)$$

$$[f_1, f_2](x) = \nabla f_2(x) f_1(x) - \nabla f_1(x) f_2(x)$$



Lie brackets: formal non-reciprocity


$$\dot{x} = \sum_{i=1}^m f_i(x) u_i$$

$\searrow m = 2$

$$\dot{x} = f_1(x)u_1 + f_2(x)u_2$$

$$x(4\varepsilon) = x_0 + \varepsilon^2[f_1, f_2](x_0) + o(\varepsilon^2)$$

$$[f_1, f_2](x) = \nabla f_2(x)f_1(x) - \nabla f_1(x)f_2(x)$$



→ We can generate new reachable directions given by the Lie brackets of the f_i

→ of course, this can be iterated: $[[f_0, f_1], [f_0, [f_1, f_2]]]$, etc.

Conditions for controllability

$$\dot{x} = \sum_{i=1}^m f_i(x)u_i$$

Definition

Let $x_0 \in \mathbb{R}^n$. Let $\text{Lie}(f_0, \dots, f_m)$ be the set with all the iterated Lie brackets:

$$\text{Lie}(f_0, \dots, f_m) = \{f_0, \dots, f_m, [f_0, f_1], \dots, [f_0, f_m], \dots, [[f_0, [f_0, f_i]], [f_j, f_k]], \dots\}$$

Lie Algebra Rank Condition in the point x_0 :

$$\text{Span}\{g(x_0), g \in \text{Lie}(f_1, \dots, f_m)\} = \mathbb{R}^n.$$

→ “The LARC is satisfied if Lie brackets generate all possible directions”

Conditions for controllability

$$\dot{x} = \sum_{i=1}^m f_i(x)u_i$$

Definition

Let $x_0 \in \mathbb{R}^n$. Let $\text{Lie}(f_0, \dots, f_m)$ be the set with all the iterated Lie brackets:

$$\text{Lie}(f_0, \dots, f_m) = \{f_0, \dots, f_m, [f_0, f_1], \dots, [f_0, f_m], \dots, [[f_0, [f_0, f_i]], [f_j, f_k]], \dots\}$$

Lie Algebra Rank Condition in the point x_0 :

$$\text{Span}\{g(x_0), g \in \text{Lie}(f_1, \dots, f_m)\} = \mathbb{R}^n.$$

→ “The LARC is satisfied if Lie brackets generate all possible directions”

Theorem (Rashevski-Chow)

Lie Algebra Rank Condition at $x_0 \Leftrightarrow$ local controllability at x_0

LARC for all x_0 in $\mathbb{R}^n \Leftrightarrow$ controllable everywhere in time T , for all T

Conditions for controllability

$$\dot{x} = \sum_{i=1}^m f_i(x)u_i$$

Definition

Let $x_0 \in \mathbb{R}^n$. Let $\text{Lie}(f_0, \dots, f_m)$ be the set with all the iterated Lie brackets:

$$\text{Lie}(f_0, \dots, f_m) = \{f_0, \dots, f_m, [f_0, f_1], \dots, [f_0, f_m], \dots, [[f_0, [f_0, f_i]], [f_j, f_k]], \dots\}$$

Lie Algebra Rank Condition in the point x_0 :

$$\text{Span}\{g(x_0), g \in \text{Lie}(f_1, \dots, f_m)\} = \mathbb{R}^n.$$

→ “The LARC is satisfied if Lie brackets generate all possible directions”

Theorem (Rashevski-Chow)

Lie Algebra Rank Condition at $x_0 \Leftrightarrow$ local controllability at x_0

LARC for all x_0 in $\mathbb{R}^n \Leftrightarrow$ **controllable everywhere** in time T , for all T

→ **Scallop theorem** : $m = 1...$

Controllability of microswimmers: an overview

rigid motion

hydrodynamics

deformation


$$\dot{\mathcal{X}} = \mathbf{N}(\mathcal{S})\dot{\mathcal{S}}$$

The swimming equation is a **driftless control-affine system**

→ **Rashveski-Chow theorem**

→ **many results of controllability**

flagellar motion



Giraldi et al., CDC, 2013

Marchello et al. Acta Appl. Math., 2022

Controllability of microswimmers: an overview

rigid motion

hydrodynamics

deformation



$$\dot{\mathcal{X}} = \mathbf{N}(\mathcal{S})\dot{\mathcal{S}}$$

The swimming equation is a **driftless control-affine system**

→ **Rashveski-Chow theorem**

→ **many results of controllability**

flagellar motion



Controllability of microswimmers: an overview

rigid motion

hydrodynamics

deformation

$$\dot{\mathcal{X}} = \mathbf{N}(\mathcal{S})\dot{\mathcal{S}}$$


The swimming equation is a **driftless control-affine system**

→ **Rashveski-Chow theorem**


→ **many results of controllability**

minimal swimmers

Alouges et al., Discrete Contin. Dyn. Syst. 2013



flagellar motion



Giraldi et al., CDC, 2013

Marchello et al. Acta Appl. Math., 2022

Controllability of microswimmers: an overview

rigid motion

hydrodynamics

deformation

$$\dot{\mathcal{X}} = \mathbf{N}(\mathcal{S})\dot{\mathcal{S}}$$

The swimming equation is a **driftless control-affine system**

→ **Rashveski-Chow theorem**

→ **many results of controllability**


minimal swimmers

Alouges et al., Discrete Contin. Dyn. Syst. 2013

flagellar motion




ciliate motion





euglenoid motion




Lohéac, Munnier, ESAIM:COCV, 2014



Giraldi et al., CDC, 2013
Marchello et al. Acta Appl. Math., 2022



Lohéac, Takahashi, ESAIM:COCV, 2020



Controllability of microswimmers: an overview

rigid motion hydrodynamics deformation

$$\dot{\mathcal{X}} = \mathbf{N}(\mathcal{S})\dot{\mathcal{S}}$$


The swimming equation is a **driftless control-affine system**
→ **Rashveski-Chow theorem**
→ **many results of controllability**

minimal swimmers

Alouges et al., Discrete Contin. Dyn. Syst. 2013


flagellar motion

euglenoid motion



NB: here, more d.o.f. makes it « easier »!

ier, ESAIM:COCV, 2014



Beyond the canonical swimming model

Questionable assumptions

1. Newtonian flow

2. Free-space swimming

3. Quiescent fluid

4. No external effects

5. Control by deformation

Beyond the canonical swimming model

$$\dot{\mathcal{X}} = \cancel{\mathbf{N}(\mathcal{S})} \dot{\mathcal{S}}$$

1. Non-Newtonian flow

$$\longrightarrow \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}, t) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$

Memory effect

or non-linear, more realistically

2. Free-space swimming

▪ Virtually nothing on controllability

3. Quiescent fluid

4. No external effects

5. Control by deformation

Beyond the canonical swimming model

$$\dot{\mathcal{X}} = \cancel{\mathbf{N}(\mathcal{S})} \dot{\mathcal{S}}$$

1. Non-Newtonian flow

$$\longrightarrow \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}, t) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$

Memory effect

2. Obstacles

$$\longrightarrow \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}, \mathcal{X}) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$


Dependence
in space

3. Quiescent fluid

4. No external effects

5. Control by deformation

- **Near a wall:** Chambrion & Munnier 2010; Alouges & Giraldi 2013
- **Multiple swimmers:** Walker, M. et al. 2022, Zoppello, Shum et al. 2024



Beyond the canonical swimming model

$$\dot{\mathcal{X}} = \cancel{\mathbf{N}(\mathcal{S})} \dot{\mathcal{S}}$$

1. Non-Newtonian flow

$$\longrightarrow \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}, t) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$

Memory effect

2. Obstacles

$$\longrightarrow \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}, \mathcal{X}) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$

Dependence
in space

3. Background flow

$$\begin{array}{l} \searrow \\ \nearrow \end{array} \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}} + \mathbf{F}_{\text{ext}}(\mathcal{X}, \mathcal{S})$$

Drift


4. Gravity, magnetism

5. Control by deformation


Beyond the canonical swimming model

$$\dot{\mathcal{X}} = \cancel{\mathbf{N}(\mathcal{S})} \dot{\mathcal{S}}$$


1.



2.



3.



$$\begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}, t) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$

Memory effect

$$\begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}, \mathcal{X}) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$


Dependence
in space

$$\begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}} + \mathbf{F}_{\text{ext}}(\mathcal{X}, \mathcal{S})$$

Drift

4. Gravity, mag

5. Control by c



- Background flow as control

Beyond the canonical swimming model

$$\dot{\mathcal{X}} = \cancel{\mathbf{N}(\mathcal{S})} \dot{\mathcal{S}}$$

1. Non-Newtonian flow

$$\longrightarrow \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}, t) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$

Memory effect

2. Obstacles

$$\longrightarrow \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}, \mathcal{X}) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$

Dependence
in space

3. Background flow

$$\begin{matrix} \searrow \\ \nearrow \end{matrix} \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}} + \mathbf{F}_{\text{ext}}(\mathcal{X}, \mathcal{S})$$

Drift

4. Gravity, magnetism

5. Indirect control

$$\longrightarrow \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}) \dot{\mathcal{S}} \\ \mathbf{Q}_0(\mathcal{S}) \end{pmatrix} + \begin{pmatrix} 0 \\ \mathbf{Q}(\mathcal{S}) \end{pmatrix} \mathcal{U}$$

Drift &
bigger state

Beyond the canonical swimming model

$$\dot{\mathcal{X}} = \cancel{\mathbf{N}(\mathcal{S})} \dot{\mathcal{S}}$$

1. Non-Newtonian flow

$$\longrightarrow \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}, t) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$

Memory effect

2. Obstacles

$$\longrightarrow \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}, \mathcal{X}) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}}$$

Dependence
in space

3. Background flow

$$\begin{matrix} \searrow \\ \nearrow \end{matrix} \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}) \\ \text{Id} \end{pmatrix} \dot{\mathcal{S}} +$$

- **Elasticity:** Alouges, DeSimone, Giraldi, Or, Zoppello, etc.
- **Magneto-elastic:** see later


4. Gravity, magnetism

5. Indirect control


$$\longrightarrow \begin{pmatrix} \dot{\mathcal{X}} \\ \dot{\mathcal{S}} \end{pmatrix} = \begin{pmatrix} \mathbf{N}(\mathcal{S}) \dot{\mathcal{S}} \\ \mathbf{Q}_0(\mathcal{S}) \end{pmatrix} + \begin{pmatrix} 0 \\ \mathbf{Q}(\mathcal{S}) \end{pmatrix} \mathcal{U}$$

**Drift &
bigger state**

A case with multiple « swimmers »



A case with multiple « swimmers »



- Motivation: control of **multiple swimmers / cargo transport**
- Two spherical particles
- A force $\mathbf{f}$ is applied on one of them
- Control of the position of **both at the same time**


- State : $(\mathbf{x}_1, \mathbf{r})$ (6-dimensional), control : $\mathbf{f} = (f_x, f_y, f_z)$

“Swimming equation” in Stokes flow :

$$\frac{d}{dt} \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{r} \end{bmatrix} = \begin{bmatrix} M_{\parallel}(r) \frac{\mathbf{r}\mathbf{r}^T}{r^2} + M_{\perp}(r) \left(\mathbf{I} - \frac{\mathbf{r}\mathbf{r}^T}{r^2} \right) \\ -m_{\parallel}(r) \frac{\mathbf{r}\mathbf{r}^T}{r^2} - m_{\perp}(r) \left(\mathbf{I} - \frac{\mathbf{r}\mathbf{r}^T}{r^2} \right) \end{bmatrix} \mathbf{f} = f_x \mathbf{g}_x + f_y \mathbf{g}_y + f_z \mathbf{g}_z$$

- **Driftless control system** $\rightarrow$ Check the LARC for the vector fields $\mathbf{g}_x, \mathbf{g}_y, \mathbf{g}_z$

A case with multiple « swimmers »



- Consider the matrix

$$C = [\mathbf{g}_x, \mathbf{g}_y, \mathbf{g}_x, [\mathbf{g}_x, \mathbf{g}_y], [\mathbf{g}_x, \mathbf{g}_z], [\mathbf{g}_y, [\mathbf{g}_x, \mathbf{g}_y]]]$$

- Its determinant at r is given by

$$\frac{m_{\parallel}^4 m_{\perp}^6}{r} \left(\frac{1}{r} \left(\frac{M_{\parallel}}{m_{\parallel}} - \frac{M_{\perp}}{m_{\perp}} \right) - \left(\frac{M_{\perp}}{m_{\perp}} \right)' \right)^3$$


Proposition (Walker, Ishimoto, Gaffney, M., JFM 2022)

- The system is controllable at r if (and only if)

$$\frac{1}{r} \left(\frac{M_{\parallel}}{m_{\parallel}} - \frac{M_{\perp}}{m_{\perp}} \right) - \left(\frac{M_{\perp}}{m_{\perp}} \right)' \neq 0.$$


A case with multiple « swimmers »

- The mobility coefficients $M_{\parallel}, m_{\parallel}, M_{\perp}, m_{\perp}$, can be approximated with series in powers of $\frac{1}{r}$
- Numerical estimation of the determinant




- controllable everywhere!
- more particles...?

- **Motion planning:** explicit build a trajectory, using the Lie brackets as “basis vectors”



Leaving the driftless territory...



El Alaoui-Faris et al. 2019

Conditions for controllability: with a drift

$$(\text{Aff}) : \dot{x} = f_0(x) + \sum_{i=1}^m f_i(x)u_i$$

- More complicated when $f_0 \neq 0$: the LARC is only necessary. Counterexample:

$$\dot{x}_1 = x_2^2, \dot{x}_2 = u$$

- **Heuristics:** brackets with an even number of times f_0 are « bad » ...

No NS condition for STLC is known!

Scalar-input case

$$\dot{x} = f_0(x) + f_1(x)u$$

- **Sufficient conditions:** Sussmann 1986

$$\text{LARC} \ \& \ (\forall k, S_{2k} \subset S_{2k-1})$$

- **Necessary conditions:** Sussmann, Kawski, Krastanov, Beauchard, Marbach, etc.

$$S_2 \not\subset S_1?$$

$$[f_1, [f_0, f_1]]$$

Two-control case

$$\dot{x} = f_0(x) + f_1(x)u_1 + f_2(x)u_2$$

- **Sufficient condition:** Sussmann « $S(\theta)$ »
- **Necessary conditions:** none!
 - 2024: one result coming from a micro-swimmer model!
 - Update 2025: now additional necessary conditions! (Gherdaoui)

Two-control case

$$\dot{x} = f_0(x) + f_1(x)u_1 + f_2(x)u_2$$

- **Sufficient condition:** Sussmann « $S(\theta)$ »
- **Necessary conditions:** none!
 - 2024: one result coming from a micro-swimmer model!
 - Update 2025: now additional necessary conditions! (Gherdaoui)

Theorem (Giraldi, Lissy, M., Pomet, ESAIM:COCV 2024)

Assume $f_2(0) = 0$ and $f_{101}(0) \notin R_1$.

1. If $f_{101}(0) \in R_1 + \text{Span}(f_{121}(0))$ let $\beta \in \mathbb{R}$ such that $f_{101}(0) + \beta f_{121}(0) \in R_1$.

Then, for all $u_{\text{eq},2} \in \mathbb{R}$ such that $u_{\text{eq},2} \neq \beta$, the system is **not STLC** at $(0, (0, u_{\text{eq},2}))$.

2. If $f_{101}(0) \notin R_1 + \text{Span}(f_{121}(0))$, then, for all $u_{\text{eq},2} \in \mathbb{R}$, the system is **not B-STLC** at $(0, (0, u_{\text{eq},2}))$.

- **One may neutralize the bad bracket f_{101} with f_{121}**

Two-control case

$$\dot{x} = f_0(x) + f_1(x)u_1 + f_2(x)u_2$$

- **Sufficient condition:** Sussmann « $S(\theta)$ »
- **Necessary conditions:** none!
 - 2024: one result coming from a micro-swimmer model!
 - Update 2025: now additional necessary conditions! (Gherdaoui)

Theorem (Giraldi, Lissy, M., Pomet, ESAIM:COCV 2024)

Assume $f_2(0) = 0$ and $f_{101}(0) \notin R_1$.



1. If $f_{101}(0) \in R_1 + \text{Span}(f_{121}(0))$ let $\beta \in \mathbb{R}$ such that $f_{101}(0) + \beta f_{121}(0) \in R_1$.

Then, for all $u_{\text{eq},2} \in \mathbb{R}$ such that $u_{\text{eq},2} \neq \beta$, the system is **not STLC** at $(0, (0, u_{\text{eq},2}))$.


2. If $f_{101}(0) \notin R_1 + \text{Span}(f_{121}(0))$, then, for all $u_{\text{eq},2} \in \mathbb{R}$, the system is **not B-STLC** at $(0, (0, u_{\text{eq},2}))$.

- One may neutralize the bad bracket f_{101} with f_{121}

Drift system: **magneto-elastic** Purcell swimmer




Drift system: **magneto-elastic** Purcell swimmer



- State: $\mathbf{z} = (x, y, \theta, \alpha_1, \alpha_2)$
- Controls: $(H_{\parallel}, H_{\perp})$

Drift system: **magneto-elastic** Purcell swimmer




$$\left\{ \begin{array}{lcl} \mathbf{f}_1^h + \mathbf{f}_2^h + \mathbf{f}_3^h & = & 0 \\ m_1^h + m_2^h + m_3^h & = & m_1^{mag} + m_2^{mag} + m_3^{mag} \\ m_2^h + m_3^h & = & m_1^{el} + m_2^{mag} + m_3^{mag} \\ m_3^h & = & m_2^{el} + m_3^{mag} \end{array} \right.$$

Equations of motion

- State: $\mathbf{z} = (x, y, \theta, \alpha_1, \alpha_2)$
- Controls: $(H_{\parallel}, H_{\perp})$

Drift system: **magneto-elastic** Purcell swimmer




$$\left\{ \begin{array}{l} \mathbf{f}_1^h + \mathbf{f}_2^h + \mathbf{f}_3^h \\ m_1^h + m_2^h + m_3^h \\ m_2^h + m_3^h \\ m_3^h \end{array} \right. = \begin{array}{l} 0 \\ \\ \\ \end{array} = \begin{array}{l} \\ \\ m_1^{el} + m_2^{mag} + m_3^{mag} \\ m_2^{el} + m_3^{mag} \end{array}$$

Hydrodynamics
Elasticity
Magnetism

Equations of motion

- State: $\mathbf{z} = (x, y, \theta, \alpha_1, \alpha_2)$
- Controls: $(H_{\parallel}, H_{\perp})$

Drift system: **magneto-elastic** Purcell swimmer




$$\left\{ \begin{array}{l} \mathbf{f}_1^h + \mathbf{f}_2^h + \mathbf{f}_3^h = 0 \\ m_1^h + m_2^h + m_3^h = m_1^{el} + m_2^{mag} + m_3^{mag} \\ m_2^h + m_3^h = m_2^{el} + m_3^{mag} \\ m_3^h = m_3^{mag} \end{array} \right.$$

$$m_i^{mag} = M_i \mathbf{H} \times \mathbf{e}_{i,\parallel}$$

- State: $\mathbf{z} = (x, y, \theta, \alpha_1, \alpha_2)$
- Controls: $(H_{\parallel}, H_{\perp})$

Drift system: **magneto-elastic** Purcell swimmer




$$\left\{ \begin{array}{lcl} \mathbf{f}_1^h + \mathbf{f}_2^h + \mathbf{f}_3^h & = & 0 \\ m_1^h + m_2^h + m_3^h & = & m_1^{el} + m_1^{mag} + m_2^{mag} + m_3^{mag} \\ m_2^h + m_3^h & = & m_2^{el} + m_2^{mag} + m_3^{mag} \\ m_3^h & = & m_3^{el} + m_3^{mag} \end{array} \right.$$

$$m_i^{mag} = M_i \mathbf{H} \times \mathbf{e}_{i,\parallel}$$

- State: $\mathbf{z} = (x, y, \theta, \alpha_1, \alpha_2)$
- Controls: $(H_{\parallel}, H_{\perp})$
- Equilibrium points: $\alpha_1 = \alpha_2 = 0, H_{\perp} = 0, H_{\parallel}$ **arbitrary**

Drift system: **magneto-elastic** Purcell swimmer



$$\left\{ \begin{array}{lcl} \mathbf{f}_1^h + \mathbf{f}_2^h + \mathbf{f}_3^h & = & 0 \\ m_1^h + m_2^h + m_3^h & = & m_1^{mag} + m_2^{mag} + m_3^{mag} \\ m_2^h + m_3^h & = & m_1^{el} + m_2^{mag} + m_3^{mag} \\ m_3^h & = & m_2^{el} + m_3^{mag} \end{array} \right.$$


$$m_i^{mag} = M_i \mathbf{H} \times \mathbf{e}_{i,\parallel}$$

$$\dot{\mathbf{z}} = \mathbf{F}_0(\alpha_1, \alpha_2) + \mathbf{F}_{\parallel}(\alpha_1, \alpha_2)H_{\parallel}(t) + \mathbf{F}_{\perp}(\alpha_1, \alpha_2)H_{\perp}(t)$$

Control-affine system with a drift

- State: $\mathbf{z} = (x, y, \theta, \alpha_1, \alpha_2)$
- Controls: $(H_{\parallel}, H_{\perp})$
- Equilibrium points: $\alpha_1 = \alpha_2 = 0, H_{\perp} = 0, H_{\parallel}$ **arbitrary**

Drift system: **magneto-elastic** Purcell swimmer



$$\left\{ \begin{array}{lcl} \mathbf{f}_1^h + \mathbf{f}_2^h + \mathbf{f}_3^h & = & 0 \\ m_1^h + m_2^h + m_3^h & = & m_1^{mag} + m_2^{mag} + m_3^{mag} \\ m_2^h + m_3^h & = & m_1^{el} + m_2^{mag} + m_3^{mag} \\ m_3^h & = & m_2^{el} + m_3^{mag} \end{array} \right.$$


$$m_i^{mag} = M_i \mathbf{H} \times \mathbf{e}_{i,\parallel}$$

$$\dot{\mathbf{z}} = \mathbf{F}_0(\alpha_1, \alpha_2) + \mathbf{F}_{\parallel}(\alpha_1, \alpha_2)H_{\parallel}(t) + \mathbf{F}_{\perp}(\alpha_1, \alpha_2)H_{\perp}(t)$$

Around which points (= **which value of $H_{\parallel}$**)
do we have local controllability?

- State: $\mathbf{z} = (x, y, \theta, \alpha_1, \alpha_2)$
- Controls: $(H_{\parallel}, H_{\perp})$
- Equilibrium points: $\alpha_1 = \alpha_2 = 0, H_{\perp} = 0, H_{\parallel}$ **arbitrary**

Drift system: **magneto-elastic** Purcell swimmer




$$\left\{ \begin{array}{lcl} \mathbf{f}_1^h + \mathbf{f}_2^h + \mathbf{f}_3^h & = & 0 \\ m_1^h + m_2^h + m_3^h & = & m_1^{mag} + m_2^{mag} + m_3^{mag} \\ m_2^h + m_3^h & = & m_1^{el} + m_2^{mag} + m_3^{mag} \\ m_3^h & = & m_2^{el} + m_3^{mag} \end{array} \right.$$

$$\dot{\mathbf{z}} = \mathbf{F}_0(\alpha_1, \alpha_2) + \mathbf{F}_{\parallel}(\alpha_1, \alpha_2)H_{\parallel}(t) + \mathbf{F}_{\perp}(\alpha_1, \alpha_2)H_{\perp}(t)$$

Proposition: The swimmer is locally controllable if and only if

$$H_{\parallel} = \frac{[\mathbf{F}_{\perp}, [\mathbf{F}_0, \mathbf{F}_{\perp}]](0,0)}{[\mathbf{F}_{\perp}, [\mathbf{F}_{\parallel}, \mathbf{F}_{\perp}]](0,0)}.$$

Drift system: **magneto-elastic** Purcell swimmer




$$\left\{ \begin{array}{lcl} \mathbf{f}_1^h + \mathbf{f}_2^h + \mathbf{f}_3^h & = & 0 \\ m_1^h + m_2^h + m_3^h & = & m_1^{mag} + m_2^{mag} + m_3^{mag} \\ m_2^h + m_3^h & = & m_1^{el} + m_2^{mag} + m_3^{mag} \\ m_3^h & = & m_2^{el} + m_3^{mag} \end{array} \right.$$

$$\dot{\mathbf{z}} = \mathbf{F}_0(\alpha_1, \alpha_2) + \mathbf{F}_{\parallel}(\alpha_1, \alpha_2)H_{\parallel}(t) + \mathbf{F}_{\perp}(\alpha_1, \alpha_2)H_{\perp}(t)$$

Proposition: The swimmer is STLC if and only if

$$H_{\parallel} = \frac{[\mathbf{F}_{\perp}, [\mathbf{F}_0, \mathbf{F}_{\perp}]](0,0)}{[\mathbf{F}_{\perp}, [\mathbf{F}_{\parallel}, \mathbf{F}_{\perp}]](0,0)} = \kappa \frac{17(M_1 + M_3) - 16M_2}{-7M_2^2 + 9M_2(M_1 + M_3) + 5M_1M_3}.$$


Drift system: **magneto-elastic** Purcell swimmer




Proposition: The swimmer is STLC if and only if

$$H_{\parallel} = \frac{[\mathbf{F}_{\perp}, [\mathbf{F}_0, \mathbf{F}_{\perp}]](0,0)}{[\mathbf{F}_{\perp}, [\mathbf{F}_{\parallel}, \mathbf{F}_{\perp}]](0,0)} = \kappa \frac{17(M_1 + M_3) - 16M_2}{-7M_2^2 + 9M_2(M_1 + M_3) + 5M_1M_3}.$$



“The control $H_{\parallel}$ must be close to this value in order to neutralise the drift”




$$H_{\parallel} < \beta$$




$$H_{\parallel} = \beta$$



$$H_{\parallel} > \beta$$






Mystery solved




El Alaoui-Faris et al. 2019

Magnetic field: vertical oscillating
+ constant horizontal




Mystery solved





El Alaoui-Faris et al. 2019



Magnetic field: vertical oscillating
+ constant horizontal


$$H_{\parallel} < \frac{[\mathbf{F}_{\perp}, [\mathbf{F}_0, \mathbf{F}_{\perp}]](0,0)}{[\mathbf{F}_{\perp}, [\mathbf{F}_{\parallel}, \mathbf{F}_{\perp}]](0,0)}$$


$$H_{\parallel} = \frac{[\mathbf{F}_{\perp}, [\mathbf{F}_0, \mathbf{F}_{\perp}]](0,0)}{[\mathbf{F}_{\perp}, [\mathbf{F}_{\parallel}, \mathbf{F}_{\perp}]](0,0)}$$


$$H_{\parallel} > \frac{[\mathbf{F}_{\perp}, [\mathbf{F}_0, \mathbf{F}_{\perp}]](0,0)}{[\mathbf{F}_{\perp}, [\mathbf{F}_{\parallel}, \mathbf{F}_{\perp}]](0,0)}$$

Two-control case, higher-order brackets

- ▶ seven other brackets of order 5 are involved, via the map

$$D_{u_2^{\text{eq}}}(\lambda_1, \lambda_2) = \lambda_1^2(f_{01,001}(0) - u_2^{\text{eq}}f_{01,201}(0)) + \lambda_2^2(f_{21,021}(0) - u_2^{\text{eq}}f_{21,221}(0)) \\ - \lambda_1\lambda_2(f_{21,001}(0) + f_{01,021}(0) - u_2^{\text{eq}}(f_{21,201}(0) + f_{01,221}(0))),$$

Theorem

Assume that $f_{101}(0)$, $f_{121}(0)$ and $f_{21,01}(0)$ belong to R_1 and let $u_2^{\text{eq}} \in \mathbb{R}$. Then :

1. If there exists a linear form φ on $\mathbb{R}^n$ such that

$$R_1 \subset \ker \varphi, \quad (12)$$

$$(\lambda_1, \lambda_2) \mapsto \langle \varphi, D_{u_2^{\text{eq}}}(\lambda_1, \lambda_2) \rangle \text{ is positive definite,} \quad (13)$$

then system (S_2) is not $(W^{1,\infty}, L^\infty)$ -STLC at $(0, (0, u_2^{\text{eq}}))$.

2. If there exists a linear form ψ on $\mathbb{R}^n$ such that

$$R' \subset \ker \psi, \quad (14)$$

$$(\lambda_1, \lambda_2) \mapsto \langle \psi, D_{u_2^{\text{eq}}}(\lambda_1, \lambda_2) \rangle \text{ is positive definite,} \quad (15)$$

then, system (S_2) is not $(W^{1,\infty}, B)$ -STLC at $(0, (0, u_2^{\text{eq}}))$.

$$[[f_0, f_1], [f_0, [f_0, f_1]]]$$

$$[[f_2, f_1], [f_0, [f_0, f_1]]]$$

...





$$[[f_2, f_1], [f_2, [f_2, f_1]]]$$

- ▶ structurally similar to the one for f_{101} but more technical

Summary

- Overview of control theory applied to microswimming
- Future challenges:
 - More controllability results for systems with a drift?
 - Multiple swimmers problem : controllability for more particles
 - “Break more assumptions” from the canonical swimming equation

References



Thank you for your attention!